

On the stability of J^* -derivations

¹M. Eshaghi Gordji, ²S. Shams, ³A. Ebadian and ⁴M. B. Ghaemi

¹Department of Mathematics, Semnan University,
P. O. Box 35195-363, Semnan, Iran

e-mail: madjid.eshaghi@gmail.com

^{2,3}Department of Mathematics, Urmia University, Urmia, Iran
e-mail: ebadian.a@gmail.com, Sa40shams@yahoo.com

⁴Department of Mathematics, Iran University of Science and Technology,
Narmak, Tehran, Iran
e-mail: mghaemi@iust.ac.ir

Abstract. In this paper, we establish the stability and superstability of J^* -derivations in J^* -algebras for the generalized Jensen-type functional equation

$$rf\left(\frac{x+y}{r}\right) + rf\left(\frac{x-y}{r}\right) = 2f(x).$$

Finally, we investigate the stability of J^* -derivations by using the fixed point alternative.

1. INTRODUCTION

Let $\mathcal{H}, \mathcal{K}$ be two Hilbert spaces and let $\mathcal{B}(\mathcal{H}, \mathcal{K})$ be the space of all bounded operators from $\mathcal{H}$ into $\mathcal{K}$. By a J^* -algebra we mean a closed subspace $\mathcal{A}$ of $\mathcal{B}(\mathcal{H}, \mathcal{K})$ such that $xx^*x \in \mathcal{A}$ whenever $x \in \mathcal{A}$. Many familiar spaces are J^* -algebras [7]. Of course J^* -algebras are not algebras in the ordinary sense. However from the point of view they may be considered a generalization of C^* -algebras; see [6, 7, 8]. In particular any Hilbert space may be thought of as a J^* -algebra identified with $\mathcal{L}(\mathcal{H}, \mathbb{C})$. Also any C^* -algebra in $\mathcal{B}(\mathcal{H})$ is a J^* -algebra. Other important examples of J^* -algebras are the so-called Cartan factors of type I, II, III and IV. A J^* -derivation on a J^* -algebra $\mathcal{A}$ is defined to be a linear mapping $d : \mathcal{A} \rightarrow \mathcal{A}$ such that

$$d(aa^*a) = d(a)a^*a + a(d(a))^*a + aa^*d(a)$$

for all $a \in \mathcal{A}$.

In particular, every $*$ -derivation on a C^* -algebra is a J^* -derivation.

The stability of functional equations was first introduced by S. M. Ulam [21] in 1940. More precisely, he proposed the following problem: Given a group G_1 , a metric group (G_2, d) and a positive number ϵ , does there exist a $\delta > 0$ such that if a function $f : G_1 \rightarrow G_2$ satisfies the inequality $d(f(xy), f(x)f(y)) < \delta$ for all $x, y \in G_1$, then there exists a homomorphism $T : G_1 \rightarrow G_2$ such that $d(f(x), T(x)) < \epsilon$ for all $x \in G_1$? As mentioned above, when this problem has a solution, we say that the homomorphisms from G_1 to G_2 are stable. In 1941, D. H. Hyers [9] gave a partial solution of *Ulam's* problem for the case of approximate additive mappings under the assumption that G_1 and G_2 are Banach spaces. In 1978, Th. M. Rassias [18] generalized the theorem of Hyers by considering the stability problem with unbounded Cauchy differences. This phenomenon of stability that was introduced by Th.

⁰2000 Mathematics Subject Classification. Primary 39B82; secondary 47B48; 46L05; 39B52; 46K99; 16Wxx.

⁰Keywords: Hyers–Ulam–Rassias stability; J^* -algebra.

M. Rassias [18] is called the Hyers–Ulam–Rassias stability. According to Th. M. Rassias theorem:

Theorem 1.1. *Let $f : E \longrightarrow E'$ be a mapping from a norm vector space E into a Banach space E' subject to the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p)$$

for all $x, y \in E$, where ϵ and p are constants with $\epsilon > 0$ and $p < 1$. Then there exists a unique additive mapping $T : E \longrightarrow E'$ such that

$$\|f(x) - T(x)\| \leq \frac{2\epsilon}{2-2^p} \|x\|^p$$

for all $x \in E$. If $p < 0$ then inequality (1.3) holds for all $x, y \neq 0$, and (1.4) for $x \neq 0$. Also, if the function $t \mapsto f(tx)$ from $\mathbb{R}$ into E' is continuous for each fixed $x \in E$, then T is linear.

During the last decades several stability problems of functional equations have been investigated by many mathematicians. A large list of references concerning the stability of functional equations can be found in [5, 10, 11, 12, 19].

Recently, Cădariu and Radu applied the fixed point method to the investigation of the functional equations. (see also [2, 3, 4, 16, 17, 20]). In [14], Park establish the stability of homomorphisms between C^* -algebras (see also [13, 15, 16]). In section 2 of the present paper, we establish the stability and superstability of J^* -derivations in J^* -algebras for the generalized Jensen-type functional equation

$$rf\left(\frac{x+y}{r}\right) + rf\left(\frac{x-y}{r}\right) = 2f(x). \quad (1.1)$$

In section 3, we will use the fixed point alternative of Cădariu and Radu to prove the stability and superstability of J^* -derivations on J^* -algebras for the generalized Jensen-type functional equation (1.1).

Throughout this paper assume that $\mathcal{A}$ is a J^* -algebra.

2. STABILITY

We start our work by a theorem in superstability of J^* -derivations.

Theorem 2.1. *Let $r, s \in (1, \infty)$, and let $D : \mathcal{A} \rightarrow \mathcal{A}$ be a mapping for which $D(sa) = sD(a)$ for all $a \in \mathcal{A}$. Suppose there exists a function $\phi : \mathcal{A}^3 \rightarrow [0, \infty)$ such that*

$$\lim_n s^{-n} \phi(s^n a, s^n b, s^n c) = 0,$$

$$\|r\mu D\left(\frac{a+b}{r}\right) + r\mu D\left(\frac{a-b}{r}\right) - 2D(\mu a) + D(cc^*c) - D(c)(c)^*c - cD(c)^*c - cc^*D(c)\| \leq \phi(a, b, c), \quad (2.1)$$

for all $\mu \in \mathbb{T}$ and all $a, b, c \in \mathcal{A}$. Then D is a J^ -derivation.*

Proof. Put $\mu = a = b = 0$ in (2.1). Then

$$\begin{aligned} & \|D(cc^*c) - D(c)c^*c - cD(c)^*c - cc^*D(c)\| \\ &= \frac{1}{s^{3n}} \|D((s^n c)(s^n c^*)(s^n c)) - D(s^n c)(s^n c^*)(s^n c) - (s^n c)D(s^n c^*)(s^n c) \\ & \quad - (s^n c)(s^n c^*)D(s^n c)\| \leq \frac{1}{s^{3n}} \phi(0, 0, s^n c) \leq \frac{1}{s^n} \phi(0, 0, s^n c) \end{aligned}$$

for all $c \in \mathcal{A}$. The right-hand side tends to zero as $n \rightarrow \infty$. So

$$D(cc^*c) = D(c)c^*c + cD(c)^*c + cc^*D(c)$$

for all $c \in \mathcal{A}$. Similarly, one can show that

$$r\mu D\left(\frac{a+b}{r}\right) + r\mu D\left(\frac{a-b}{r}\right) = 2D(\mu a)$$

for all $\mu \in \mathbb{T}$ and all $a, b \in \mathcal{A}$. Hence, D is linear. $\square$

Theorem 2.2. *Let $r \in (1, \infty)$, and let $f : \mathcal{A} \rightarrow \mathcal{A}$ be a mapping with $f(0) = 0$ for which there exists a function $\phi : \mathcal{A}^3 \rightarrow [0, \infty)$ such that*

$$\Phi(a, b, c) := \sum_0^\infty 2^{-n} \phi(2^n a, 2^n b, 2^n c) < \infty,$$

$$\|r\mu f\left(\frac{a+b}{r}\right) + r\mu f\left(\frac{a-b}{r}\right) - 2f(\mu a) + f(cc^*c) - f(c)(c)^*c - cf(c)^*c - cc^*f(c)\| \leq \phi(a, b, c), \quad (2.2)$$

for all $\mu \in \mathbb{T}$ and all $a, b, c \in \mathcal{A}$. Then there exists a unique J^* -derivation $D : \mathcal{A} \rightarrow \mathcal{A}$ such that

$$\|f(a) - D(a)\| \leq \Phi(a, a, 0) \quad (2.3)$$

for all $a \in \mathcal{A}$.

Proof. Put $\mu = 1$ and $b = c = 0$ in (2.2). It follows that

$$\|f(a) - r^{-1}f(ra)\| \leq \frac{1}{2}\phi(ra, 0, 0) \quad (2.4)$$

for all $a \in \mathcal{A}$. By induction, we can show that

$$\|f(a) - r^{-n}f(r^n a)\| \leq \frac{1}{2} \sum_1^n \phi(r^n a, 0, 0) \quad (2.5)$$

for all $a \in \mathcal{A}$. Replace a by a^m in (2.5) and then divide by r^m , we get

$$\|f(a^m) - r^{-n-m}f(r^{n+m}a)\| \leq \frac{1}{2r^m} \sum_m^{m+n} \phi(r^k a, 0, 0)$$

for all $a \in \mathcal{A}$. Hence, $\{r^{-n}f(r^n a)\}$ is a Cauchy sequence. Since $\mathcal{A}$ is complete, then

$$D(a) := \lim_n r^{-n}f(r^n a)$$

exists for all $a \in \mathcal{A}$. By using (2.1) one can show that

$$\begin{aligned} & \|rD\left(\frac{a+b}{r}\right) + rD\left(\frac{a-b}{r}\right) - 2D(a)\| \\ &= \lim_n \frac{1}{r^n} \|rf(r^{n-1}(a+b)) + rf(r^{n-1}(a-b)) - 2f(r^n a)\| \\ &\leq \lim_n \frac{1}{r^n} \phi(r^n a, r^n b, 0) = 0 \end{aligned}$$

for all $a, b \in \mathcal{A}$. So

$$rD\left(\frac{a+b}{r}\right) + rD\left(\frac{a-b}{r}\right) = 2D(a)$$

for all $a, b \in \mathcal{A}$. Put $U = \frac{a+b}{r}$, $V = \frac{a-b}{r}$ in above equation, we get

$$r(D(U) + D(V)) = 2rD\left(\frac{U+V}{2}\right)$$

for all $U, V \in \mathcal{A}$. Hence, D is Cauchy additive. On the other hand, we have

$$\|D(\mu a) - \mu D(a)\| = \lim_n \frac{1}{r^n} \|f(\mu r^n a) - \mu f(r^n a)\| \leq \lim_n \frac{1}{r^n} \phi(r^n a, r^n a, 0) = 0$$

for all $\mu \in \mathbb{T}$, and all $a \in \mathcal{A}$. So it is easy to show that D is linear. It follows from (2.1) that

$$\begin{aligned} & \|D(cc^*c) - D(c)c^*c - cD(c^*)c - cc^*D(c)\| \\ &= \lim_n \left\| \frac{1}{r^{3n}} f((r^n c)(r^n c^*)(r^n c)) - \frac{1}{r^n} f(r^n c) \frac{r^n c^*}{r^n} \frac{r^n c}{r^n} - \frac{r^n c}{r^n} \frac{1}{r^n} f(r^n c^*) \frac{r^n c}{r^n} \right. \\ & \quad \left. - \frac{r^n c}{r^n} \frac{r^n c^*}{r^n} \frac{1}{r^n} f(r^n c) \right\| \leq \lim_n \frac{1}{r^{3n}} \phi(0, 0, r^n c) \leq \lim_n \frac{1}{r^n} \phi(0, 0, r^n c) \\ &= 0 \end{aligned}$$

for all $c \in \mathcal{A}$. Thus $D : \mathcal{A} \rightarrow \mathcal{A}$ is a J^* -derivation satisfying (2.3), as desired. $\square$

We prove the following Hyers–Ulam–Rassias stability problem for J^* -derivations on J^* -algebras.

Corollary 2.3. *Let $p \in (0, 1)$, $\theta \in [0, \infty)$ be real numbers. Suppose $f : \mathcal{A} \rightarrow \mathcal{A}$ satisfies*

$$\|r\mu f(\frac{a+b}{r}) + r\mu f(\frac{a-b}{r}) - 2f(\mu a) + f(cc^*c) - f(c)(c)^*c - cf(c)^*c - cc^*f(c)\| \leq \theta(\|a\|^p + \|b\|^p + \|c\|^p),$$

for all $\mu \in \mathbb{T}$ and all $a, b, c \in \mathcal{A}$. Then there exists a unique J^* -derivation $D : \mathcal{A} \rightarrow \mathcal{A}$ such that

$$\|f(a) - D(a)\| \leq \frac{2^p \theta}{2^{p-1} - 1} \|a\|^p$$

for all $a \in \mathcal{A}$.

Proof. Setting $\phi(a, b, c) := \theta(\|a\|^p + \|b\|^p + \|c\|^p)$ for all $a, b, c \in \mathcal{A}$, in above theorem. $\square$

3. STABILITY BY USING ALTERNATIVE FIXED POINT

Before proceeding to the main results of this section, we will state the following theorem.

Theorem 3.1. *(The alternative of fixed point [1]). Suppose that we are given a complete generalized metric space (Ω, d) and a strictly contractive mapping $T : \Omega \rightarrow \Omega$ with Lipschitz constant L . Then for each given $x \in \Omega$, either*

$$d(T^m x, T^{m+1} x) = \infty \text{ for all } m \geq 0,$$

or other exists a natural number m_0 such that

- ★ $d(T^m x, T^{m+1} x) < \infty$ for all $m \geq m_0$;
- ★ the sequence $\{T^m x\}$ is convergent to a fixed point y^* of T ;
- ★ y^* is the unique fixed point of T in the set $\Lambda = \{y \in \Omega : d(T^{m_0} x, y) < \infty\}$;
- ★ $d(y, y^*) \leq \frac{1}{1-L} d(y, Ty)$ for all $y \in \Lambda$.

Theorem 3.2. *Let $f : \mathcal{A} \rightarrow \mathcal{A}$ be a mapping for which there exists a function $\phi : \mathcal{A}^3 \rightarrow [0, \infty)$ such that*

$$\|r\mu f(\frac{a+b}{r}) + r\mu f(\frac{a-b}{r}) - 2f(\mu a) + f(cc^*c) - f(c)(c)^*c - cf(c)^*c - cc^*f(c)\| \leq \phi(a, b, c), \quad (3.1)$$

for all $\mu \in \mathbb{T}$ and all $a, b, c \in \mathcal{A}$. If there exists an $L < 1$ such that

$$\phi(a, b, c) \leq rL\phi(\frac{a}{r}, \frac{b}{r}, \frac{c}{r}) \quad (3.2)$$

for all $a, b, c \in \mathcal{A}$, then there exists a unique J^* -derivation $D : \mathcal{A} \rightarrow \mathcal{A}$ such that

$$\|f(a) - D(a)\| \leq \frac{L}{1-L} \phi(a, 0, 0) \quad (3.3)$$

for all $a \in \mathcal{A}$.

Proof. Put $\mu = 1, b = c = 0$ in (3.1) to obtain

$$\|2rf(\frac{a}{r}) - 2f(a)\| \leq \phi(a, 0, 0) \quad (3.4)$$

for all $a \in \mathcal{A}$. Hence,

$$\|\frac{1}{r}f(ra) - f(a)\| \leq \frac{1}{2r}\phi(ra, 0, 0) \leq L\phi(ra, 0, 0) \quad (3.5)$$

for all $a \in \mathcal{A}$.

Consider the set $X := \{g \mid g : A \rightarrow \mathcal{A}\}$ and introduce the generalized metric on X :

$$d(h, g) := \inf\{C \in \mathbb{R}^+ : \|g(a) - h(a)\| \leq C\phi(a, 0, 0) \forall a \in \mathcal{A}\}.$$

It is easy to show that (X, d) is complete. Now we define the linear mapping $J : X \rightarrow X$ by

$$J(h)(a) = \frac{1}{r}h(ra)$$

for all $a \in \mathcal{A}$. By Theorem 3.1 of [1],

$$d(J(g), J(h)) \leq Ld(g, h)$$

for all $g, h \in X$.

It follows from (3.5) that

$$d(f, J(f)) \leq L.$$

By Theorem 3.1, J has a unique fixed point in the set $X_1 := \{h \in X : d(f, h) < \infty\}$. Let D be the fixed point of J . D is the unique mapping with

$$D(ra) = rD(a)$$

for all $a \in \mathcal{A}$ satisfying there exists $C \in (0, \infty)$ such that

$$\|D(a) - f(a)\| \leq C\phi(a, 0, 0)$$

for all $a \in \mathcal{A}$. On the other hand we have $\lim_n d(J^n(f), D) = 0$. It follows that

$$\lim_n \frac{1}{2^n} f(2^n a) = D(a) \quad (3.6)$$

for all $a \in \mathcal{A}$. It follows from $d(f, h) \leq \frac{1}{1-L}d(f, J(f))$, that

$$d(f, h) \leq \frac{L}{1-L}.$$

This implies the inequality (3.3).

It follows from (3.2) that

$$\lim_j r^{-j} \phi(r^j a, r^j b, r^j c) = 0 \quad (3.7)$$

for all $a, b, c \in \mathcal{A}$.

By a same reasoning as Theorem 2.2, one can show that the mapping $D : \mathcal{A} \rightarrow \mathcal{A}$ is a J^* -derivation satisfying (3.3), as desired. $\square$

We prove the following Hyers–Ulam–Rassias stability problem for J^* -derivations on J^* -algebras.

Corollary 3.3. Let $p \in (0, 1), \theta \in [0, \infty)$ be real numbers. Suppose $f : A \rightarrow A$ satisfies

$$\|r\mu f(\frac{a+b}{r}) + r\mu f(\frac{a-b}{r}) - 2f(\mu a) + f(cc^*c) - f(c)(c)^*c - cf(c)^*c - cc^*f(c)\| \leq \theta(\|a\|^p + \|b\|^p + \|c\|^p),$$

for all $\mu \in \mathbb{T}$ and all $a, b, c \in A$. Then there exists a unique J^* -derivation $D : A \rightarrow A$ such that

$$\|f(a) - D(a)\| \leq \frac{2^p \theta}{2 - 2^p} \|a\|^p$$

for all $a \in A$.

Proof. Setting $\phi(a, b, c) := \theta(\|a\|^p + \|b\|^p + \|c\|^p)$ all $a, b, c \in A$. Then by $L = 2^{p-1}$, we get the desired result. $\square$

Now we establish the superstability of J^* -derivations by using the alternative of fixed point.

Theorem 3.4. Let $s > 1$, and let $f : A \rightarrow A$ be a mapping satisfying $f(sx) = sf(x)$ for all $x \in A$. Let $\phi : A^3 \rightarrow [0, \infty)$ be a mapping satisfying (3.1). If there exists an $L < 1$ such that

$$\phi(x, y, z) \leq rL\phi(\frac{x}{r}, \frac{y}{r}, \frac{z}{r})$$

for all $x, y, z \in A$, then f is a J^* -derivation.

Proof. It is similar to the proof of Theorem 2.1. $\square$

Corollary 3.5. Let $r, p \in (0, 1), \theta \in [0, \infty)$ be real numbers. Suppose $f : A \rightarrow A$ is a mapping satisfying $f(rx) = rf(x)$ for all $x \in A$. Let $\phi : A^3 \rightarrow [0, \infty)$ be a mapping satisfying (3.1). Then f is a J^* -derivation.

Proof. Setting $\phi(x, y, z) := \theta(\|x\|^p + \|y\|^p + \|z\|^p)$ all $x, y, z \in A$. Then by $L = 2^{p-1}$ in above theorem, we get the desired result. $\square$

REFERENCES

- [1] L. Cădariu and V. Radu, *On the stability of the Cauchy functional equation: a fixed point approach*, Grazer Mathematische Berichte **346** (2004), 43–52.
- [2] L. Cădariu, V. Radu, *The fixed points method for the stability of some functional equations*, Carpathian Journal of Mathematics **23** (2007), 63–72.
- [3] L. Cădariu, V. Radu, *Fixed points and the stability of quadratic functional equations*, Analele Universitatii de Vest din Timisoara **41** (2003), 25–48.
- [4] L. Cădariu and V. Radu, *Fixed points and the stability of Jensen's functional equation*, J. Inequal. Pure Appl. Math. **4** (2003), Art. ID 4.
- [5] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, 2002.
- [6] M. Elin, L. Harris, S. Reich and D. Shoikhet, *Evolution Equations and Geometric Function Theory in J^* -algebras*, J. Nonlinear Convex Anal., **3** (2002), 81–121.
- [7] L.A. Harris, *Bounded symmetric homogeneous domains in infinite-dimensional spaces*, Lecture Notes in Mathematics, vol. 364, Springer, Berlin, 1974.
- [8] L.A. Harris, *Operator Siegel domains*, Proc. Roy. Soc. Edinburgh Sect. A. **79** (1977) 137156.
- [9] D.H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [10] D.H. Hyers and Th.M. Rassias, *Approximate homomorphisms*, Aequationes Math. **44** (1992), no. 2-3, 125–153.

- [11] D.H. Hyers, G. Isac and Th.M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [12] S.-M. Jung, *Hyers–Ulam–Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, 2001.
- [13] M. S. Moslehian and C. Park, *On the stability of J^* -homomorphisms*, Nonlinear Analysis, **63** (2005), 42-48.
- [14] C. Park, *Isomorphisms between C^* -ternary algebras*, J. Math. Anal. Appl. **327** (2007), 101-115.
- [15] C. Park, *Homomorphisms between Poisson JC^* -algebras*, Bull. Braz. Math. Soc. **36** (2005) 79-97. MR2132832 (2005m:39047)
- [16] C. Park and J. M. Rassias, *Stability of the Jensen-type functional equation in C^* -algebras: a fixed point approach*, Abstract and Applied Analysis Volume 2009, Article ID 360432, 17 pages.
- [17] V. Radu, *The fixed point alternative and the stability of functional equations*, Fixed Point Theory **4** (2003), 91–96.
- [18] Th.M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978) 297–300.
- [19] Th.M. Rassias, *On the stability of functional equations and a problem of Ulam*, Acta Appl. Math. **62** (2000), no. 1, 23–130.
- [20] I.A. Rus, *Principles and Applications of Fixed Point Theory*, Ed. Dacia, Cluj-Napoca, 1979 (in Romanian).
- [21] S. M. Ulam, *Problems in Modern Mathematics*, Chapter VI, science ed. Wiley, New York, 1940.